

Historic, archived document


Do not assume content reflects current scientific knowledge, policies, or practices.

A281.9
A98
cop 2
#100

Rubber Content of

MISCELLANEOUS PLANTS

EXTRA COPY



Production Research Report No. 10

Agricultural Research Service

UNITED STATES DEPARTMENT OF AGRICULTURE

USDA, National Agricultural Library
NAL Bldg
10301 Baltimore Blvd
Beltsville, MD 20705-2351

CONTENTS

	Page
Need for new sources of crude rubber.....	1
Public reaction.....	2
Type of material analyzed.....	3
Methods of analysis.....	5
Presentation of data.....	6
Conclusions.....	24
Literature cited.....	24

In addition to the contributions by the author and other personnel of the USDA laboratory, special attention should be called to the contribution by A. V. McMullan, now with the International Cooperation Administration in Costa Rica, who supervised the laboratory and made most of the analyses. All botanical identifications were made in the Division of Plant Exploration and Introduction (now the Plant Introduction Section of the Crops Research Division). The rubber work had been under that Division prior to receiving separate status at the beginning of World War II.

Rubber Content of

MISCELLANEOUS PLANTS

By LOREN G. POLHAMUS, Collaborator

Crops Research Division, Agricultural Research Service

Need for New Sources of Crude Rubber

The United States Department of Agriculture has been interested in natural sources of crude rubber since the latter part of the 19th century. Department scientists in the opening years of this century made a detailed survey of the cultivation of the Castilla rubber tree in Mexico (2).¹ At that time they urged caution to American investors who were putting millions of dollars into highly speculative Mexican rubber plantations.

The United States Department of Commerce has also been interested for many years in sources of rubber and in its ever-growing economic importance. The strategic importance of rubber was forcibly emphasized during World War I when encircled Germany tried desperately but unsuccessfully to synthesize a satisfactory substitute for the natural rubber it no longer could import.

Following the end of that war, this country recognized that rubber had become a strategic commodity and that, with regard to its sources of supply, the position of the United States might be as precarious as that of Germany. Industrialists Harvey S. Firestone and Henry Ford joined with inventor Thomas G. Edison and the then Secretary of Commerce Herbert Hoover to inform the American public as to the growing importance of rubber and the strategic liability of depending on sources on the other side of the world. "America," they said, "should produce its own rubber."

¹ Italic numbers in parentheses refer to Literature Cited, p. 24.

Congress recognized the problem, and in 1922 it granted the Secretary of Commerce an appropriation of \$500,000 to investigate sources of crude rubber. Of this sum, \$100,000 could be allocated to the Department of Agriculture to investigate rubber production in the Western Hemisphere and the Philippine Islands. This allocation from the Department of Commerce constituted the first funds received by the Department of Agriculture that were earmarked specifically for investigating the production of rubber. This first allocation has been supplemented by direct annual appropriations since then.

The Department of Commerce made a worldwide survey of rubber production and marketing and published the results of its surveys in its Trade Promotion Series (3, 6, 13, 14, 15, 16, 18). Specialists from the Department of Agriculture assisted in the surveys made by the Department of Commerce in the Western Hemisphere and in the Philippines. The Department of Agriculture published reports by its specialists on surveys in Brazil, Dutch Guiana, and the island of Trinidad (7, 12, 17). Simultaneously with the survey of existing sources of rubber, the Department of Agriculture initiated research on rubber production in the Western Hemisphere. Tapping experiments were started on a small planting of rubber trees on the northern coast of Haiti (10). These plantings consisted of *Castilla*, *Ficus*, *Funtumia*, *Hevea*, and other rubber-bearing plants. Only the *Hevea* rubber tree was found worthy of continued study.

In the United States, experimental plantings were started at Coconut Grove, Fla., and at Bard, Calif. Many tropical rubber-bearing plants were imported and studied under cultivation in Florida. The plantings at Bard, Calif., were used principally for the study of desert rubber-bearing plants, the chief of which was the desert milkweed, *Asclepias subulata*. The work of a private company with the desert rubber-bearing shrub *Parthenium argentatum* was kept under observation, but it was not felt necessary to divert any of the limited government funds to duplicate work already underway at private expense.

Public Reaction

The speculative boom in rubber planting in Mexico during the early years of the 20th century resulted in the loss of many millions of dollars by investors in the United States. It did serve to awaken the American public to the growing importance of rubber. The appeals of Firestone, Ford, Edison, and Hoover in the 1920's quickly found an answering chord among people of every age and position in the United States. Rubber became an important category in school curricula, and government agencies were flooded with requests for information from school children and their teachers. Educational pamphlets and exhibits became important parts of the informational material supplied correspondents by the large rubber-manufacturing companies.

Men, women, and children began to think in terms of "America should produce its own rubber." They knew that most of our natural rubber was obtained from the Para rubber tree. Moreover, the impression persisted that Brazil was the chief source of our rubber for many years after the increase in plantation production in the Far East placed Brazilian production in a position of relative unimportance.

Hundreds of plants throughout the world have been used as sources of crude rubber to a lesser or greater degree. In the United States, Hall and Goodspeed (4) and Hall and Long (5) of the University of California made surveys of native plants of the West that contained rubber and reported a surprising number as having significant amounts. Every plant that had a milky juice, a gummy exudate, a sticky feeling, or just no other apparent use became suspect as a potential source of rubber.

The Department of Agriculture, having assumed responsibility for investigating the production of rubber in the Americas, became recipient of inquiries as to sources of rubber. Individuals throughout the country became interested in studying the plants around them and speculating as to their potential value as sources of rubber. From 1920 to 1945, hundreds of people went to considerable personal trouble and expense to collect and furnish the Department with plants that they hoped might contain significant quantities of rubber. Many samples were supplied to the Department direct; some were furnished through other agencies in the Government; and some were supplied through members of Congress. Wherever possible, the Department made a botanical identification of the plants and a chemical analysis to determine the rubber content. Then, it furnished the individual submitting the sample with that information, together with such additional information regarding the plant as a possible source of rubber as might already be available.

Without doubt, many of the individuals who furnished samples of suspected rubber-bearing plants to the Department for analysis were imbued with the prospect of personal gain. Others were chiefly interested in performing a patriotic service.

Type of Material Analyzed

Specimens submitted to the Department for analysis varied considerably. In most cases, the collector gathered the available plant material, wrapped it up, and mailed it to the Department. Often such material arrived in a state of decay that made identification difficult and analysis uncertain. In many cases, insufficient material was sent for definite botanical identification. In some cases, evidently an effort was made to prevent identification in the hope of obtaining exclusive information that might lead to personal gain. In some cases, ground samples were submitted; sometimes extracts; often only a gum or resin; once a sample of *Hevea* latex was submitted as coming from poinsettia. To avoid furnishing reports that might be misleading, botanical specimens

were requested of plants from which questionable gums, resins, or latex were received.

Many of the specimens were received from Department personnel engaged in other activities but interested in collecting and submitting samples of interesting looking plants that they encountered in the course of their regular activities. The Department of Agriculture has made no specific survey of possible rubber-bearing plants other than that made by Mildred M. Pladeck, who was assigned to collect samples of native species of goldenrod within a 100-mile radius of Washington, D. C. A preliminary report was published in 1933 (11). For the sake of completeness, the published material is combined with that from subsequent collections in the present report.

During World War II, the Department cooperated with the Board of Economic Warfare, the Rubber Development Corporation, and other wartime agencies engaged in determining what new sources of rubber could be found in Latin America and other still-free areas of the world. Employees of these agencies, either directly supervised by the Department of Agriculture or in cooperation with the Department, submitted samples of rubbers and other gums for chemical determination of the rubber content. In the course of their search for new sources of rubber from already recognized plants, these individuals also found other plants that might be valuable in extending knowledge of the type and range of rubber-bearing plants. These plants are included in this report.

Whenever possible, samples were subdivided into the different plant parts in order to record where the rubber was formed in the different plants. However, that was not always possible, and it was necessary to analyze a composite sample of the material submitted. Such samples could not be assumed to represent a composite of the plant from which they were collected and certainly would not be considered as representative of the species. Such specimens are designated in this report as "whole" and thus are representative of the whole sample submitted. This would mean that more than just leaves or twigs or other specific portions of the plant were analyzed. However, it might mean only leaves and twigs from a bush or even a tree. If rubber is reported in such samples, there is a fair assumption that under some conditions rubber is formed in that species. If the report is that no rubber was found in such samples, it may or may not indicate that that species does not form rubber.

No coordinated survey of American rubber-bearing plants was attempted. The samples received had been collected without regard to seasonal variation in rubber content. Many plants do not accumulate rubber during periods of active growth. Rubber accumulation occurs in periods of retarded or suspended growth in most Temperate Zone plants. Analyses included in this report were of plants collected at random when the interest of the collectors dictated. It is quite probable

that in many cases higher rubber contents would have been found if the collections had been made at more favorable seasons.

The address given by the collector is shown as the point of origin of the sample unless information included with any sample indicated that it was collected elsewhere. Inclusion of plants in these lists indicates that to the best of our knowledge the plants were growing in the State or States indicated as the origin of the individual samples. The plants may have been wild or cultivated, native or introduced.

Methods of Analysis

At first, the gravimetric method of analysis described by Hall and Goodspeed (4) was used for determining the rubber content of the specimens. This is essentially a 3-hour extraction of a 5-gram ground sample with acetone, followed by a 3-hour extraction with benzene. After drying, the benzene extract was weighed direct as rubber. Later, when tests showed that the 3-hour period was insufficient to assure complete extraction, the length of the extraction periods was increased. It also was found that adding 1 percent of trichloroacetic acid to the benzene speeded up the solution of the rubber. No record was kept of the precise details of the analysis used on these miscellaneous samples. The method of analysis currently used in routine research tests was used.

In many cases, the benzene extract did not have the characteristics of true rubber. If time permitted and the sample was of some interest, the extract would be redissolved in benzene and either be precipitated with alcohol or be treated to transform the rubber into a bromide. Formation of a bromide insoluble in 95-percent alcohol was considered proof of the identity of the extract as rubber. The benzene extracts of only a few of these miscellaneous plants were checked in this manner, however, because of the time involved.

Chemically, rubber is polyisoprene. Gutta, a second polyisoprene, is also formed in plants. Both rubber, *cis*-polyisoprene, and gutta, *trans*-polyisoprene, are soluble in benzene and insoluble in acetone. The methods of analysis used in our laboratory would not differentiate between these materials. Examining the benzene extract from each specimen and noting its physical character were standard practices. The leathery extract from *Eucommia ulmoides* was known to be gutta rather than rubber, and similar extracts from other plants were assumed to be also. This was true particularly with regard to plants belonging to the Sapotaceae. A precise determination of rubber and gutta could not be made with the facilities in our laboratory. For the purpose of general testing of miscellaneous plants, the benzene extract was designated as rubber. That designation is followed in this report.

There was a significant variation in the character of the benzene extracts of plants. Some of this variation was attributed to differences

in the molecular weights of the rubber from different plants or from different parts of the same plant. In general, rubber extracts with high molecular weight are stiffer and less sticky than those with low molecular weight. Soft, sticky extracts were also sometimes attributed to ineffective extraction of the resinous nonrubber materials by acetone or to the oxidation of the rubber during the extraction period. It was on the basis of this examination that a determination was made as to whether further examination by precipitation or bromination was desirable.

Only the analyses made in the Washington, D. C., laboratories (transferred to Beltsville, Md., in 1942) are reported herein. Additional miscellaneous analyses were made for correspondents in the research laboratories in California. Those analyses have not been segregated from the research data and are not included.

Presentation of Data

Table 1 gives the results of the analyses of miscellaneous plants tested for correspondents from the continental United States and are not included in formal research activities. Nearly half the species were of the Compositae family, with fair representation from Apocynaceae, Asclepiadaceae, and Euphorbiaceae—all known to contain important rubber-bearing plants. Most of the other families are represented by only one or two species. Table 2 summarizes the total plants tested and the number that contained rubber. Table 3 gives a tabulation of the families represented, the number of species tested in each family, the number of tests made, the number that contained rubber, and the number that showed no trace of rubber.

Table 4 shows the results of analyses of plants submitted from outside the continental limits of the United States. These samples were submitted largely by individuals with considerable knowledge of rubber production, and the botanical range of the specimens was restricted almost entirely to plant families known to contain rubber-bearing plants. A total of 54 species from 33 genera, representing 7 plant families, was tested. A tabulation by family is presented in table 5.

TABLE 1.—*Rubber content of miscellaneous native and introduced plants collected in continental United States, analyzed 1920-45*

Genus and species	Family	State where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Acacia</i> sp.	Leguminosae.	California.	Gum.	1	Percent 0	Percent	Percent
<i>Acer platanoides</i> L.	Aceraceae.	Maryland.	Leaves.	1	.17		
<i>Achokanthera spectabilis</i> (Sond.) Benth.	Apocynaceae.	No record.	do.	1	.17		
<i>Adenophora</i> sp.	Campanulaceae.	District of Columbia.	Roots.	1	.20		
<i>Aesculus glabra</i> Willd.	Aesculaceae.	Virginia.	Leaves.	1	0		
<i>Agnosma acuminata</i> (Roxb.) G. Don.	Apocynaceae.	No record.	Stems.	1	0		
<i>Ailanthus altissima</i> (Mill.) Swingle.	Simarubaceae.	North Carolina.	Buckeyes.	1	0		
<i>Allium cepa</i> L.	Liliaceae.	West Virginia.	Leaves.	1	2.03		
<i>Amsonia ciliata</i> Walt.	Apocynaceae.	Georgia.	Whole.	1	4.96		
<i>Chamaenactis</i> Walt.	do.	Massachusetts.	Extract.	1	6.97		
sp.	do.	Texas.	Whole.	1	.74		
<i>Apios americana</i> Medie.	Leguminosae.	Georgia, New Jersey, Michigan.	do.	1	.37		
<i>Aplopappus ciliatus</i> (Nutt.) DC.	Compositae.	Texas.	Whole.	4	.58		
<i>divaricatus</i> (Nutt.) Gray.	do.	Stems.	do.	1	.25	0.46	0.31
<i>heterophyllus</i> (Gray) Blake.	do.	South Carolina.	Whole.	3	.41	.56	.47
<i>tenuisectus</i> (Greene) Blake.	do.	New Mexico.	Stems.	1	1.06		
<i>Apocynum androsaemifolium</i> L.	Apocynaceae.	Arizona.	Whole.	1	.59		
<i>canadinum</i> L.	do.	New Jersey, Florida, New Hampshire.	do.	1	.26		
<i>sibiricum</i> Jaed.	do.	Colorado, Maryland, New Hampshire, and Virginia.	Leaves.	2	.45	.53	.49
sp.	do.	New Mexico.	Stems.	2	.10	.14	.12
<i>Aristolochia ringens</i> Vahl.	Aristolochiaceae.	Arkansas, California, Kentucky.	Whole.	1	.56	1.42	.78
<i>Artemisia aromatica</i> A. Nels.	Compositae.	Arizona.	Leaves.	5	.50	.23	.17
<i>bigelovii</i> Gray.	do.	No record.	Stems.	2	.32	.11	.11
<i>canadensis</i> Michx.	do.	do.	Whole.	2	.33	.56	.18
<i>dracunculoides</i> Pursh.	do.	New Mexico.	Leaves.	1	.23		
			Stems.	1	.29	.63	.46
			Leaves.	2	.20	.26	.23
			Whole.	2	.83		
			Leaves.	1	.96		
			Stems.	1	1.44		
			Whole.	1	.12		
			Stems.	1	.05		
			Whole.	1	.09		
			Stems.	1	.36		
			Leaves.	1	.02		
			Stems.	1	.51		
			Whole.	1	.24		

Genus and species	Family	State where collected	Samples		Rubber content		
			P'art	Number	Least	Most	Mean
<i>Artemisia aromatica</i> A. Nels.—Continued							
<i>filifolia</i> Torr.	Compositae.	New Mexico.	(Leaves.	1	.38		<i>Percent</i>
			(Stems.	1	.11		
<i>graphalodes</i> Nutt.	do.	Arizona.	(Leaves.	2	.13	.28	.23
			(Stems.	2	.05	.12	.06
<i>ludoviciana</i> Nutt.	do.	Georgia.	(Leaves.	1	.52		
<i>scallertiana</i> Bess.	do.	District of Columbia.	(Leaves.	1	1.84		
			(Stems.	1	1.14		
<i>tridentata</i> Nutt.	do.	Montana, New Mexico.	(Leaves.	1	.37		
			(Stems.	1	.03		
<i>wrightii</i> Gray.	do.	Arizona.	(Whole.	1	.62		
			(Leaves.	1	.39		
<i>Asclepias amplexicaulis</i> J. E. Sm.	Asclepiadaceae.	North Carolina, Oklahoma.	(Stems.	1	.09		
<i>brachystephana</i> Engelm.	do.	Texas.	(Leaves.	2	2.36	4.26	3.31
			(Stems.	2	.28	.29	.28
<i>capricornu</i> Woodson.	do.	do.	(Whole.	1	.24		
			(Leaves.	1	1.06		
<i>curassavica</i> L.	do.	Florida.	(Stems.	1	.14		
			(Roots.	1	1.03		
<i>englemanniana</i> Woodson.	do.	Texas.	(Leaves.	1	.25		
			(Stems.	1	.08		
<i>orizocarpa</i> Benth.	do.	California.	(Roots.	1	.46		
<i>erosa</i> Torr.	do.	do.	(Leaves.	1	0		
<i>galiioides</i> H. B. K.	do.	Texas.	(Stems.	1	0		
<i>humistrata</i> Walt.	do.	Florida.	(Whole.	1	1.37		
			(Leaves.	1	1.60		
<i>incarnata</i> L.	do.	Texas.	(Whole.	1	2.91		
			(Stems.	1	.12		
<i>Asclepias latifolia</i> Raf.	do.	Arizona, Texas.	(Leaves.	2	(1)	1.11	.56
			(Stems.	2	0	.21	.10
<i>lindeheimeri</i> Engelm. & Gray.	do.	Texas.	(Roots.	2	0	.13	.06
			(Leaves.	4	1.89	4.46	2.47
<i>speciosa</i> Torr.	do.	Colorado, Idaho.	(Stems.	2	.18	.46	.35
			(Roots.	2	.03	.04	.04
<i>subulata</i> Decne.	do.	California.	(Whole.	1	.23		
			(Leaves.	1	.66		
			(Stems.	2	(1)	1.32	.37
			(Gum.	1	6.07		.18
			(Leaves.	1	.25		
			(Whole.	4	3.28	4.23	3.79

<i>sullivanii</i> Engelm.	do.	Texas	Whole.	1	1.15	4.84	2.40
<i>syriaca</i> L.	do.	Delaware, Illinois, Kentucky, Massachusetts, New York, Pennsylvania, Virginia, District of Columbia.	Stems.	8	1.53	5.50	2.42
<i>viridiflora</i> Raf.	do.	Oklahoma.	Whole.	3	.52	2.26	1.15
sp.	do.		Leaves.	1	1.57		
<i>Asclepiodora nudiflora</i> (Walt.) Gray.	do.	Idaho, Missouri.	Stems.	1	1.15		
<i>Asclepiodora maculata</i> (Turner) Holmes & Batters.	do.	No record.	Latex.	1	7.54		
<i>Aster pauciflorus</i> L.	Fucaceae.	Connecticut.	Whole.	1	.25		
<i>reticulatus</i> Pursh.	Compositae.	Maryland.	Whole.	1	.67		
<i>spinosus</i> Benth.	do.	South Carolina.	do.	2	0		
sp.	do.	Texas.	Leaves.	1	1.61	3.57	2.59
<i>Baccharis glutinosa</i> Pers.	do.	Georgia.	Stems.	1	.34		
<i>ladimifolia</i> L.	do.	Texas.	Roots.	1	.06		
<i>neglecta</i> Britton.	do.	Mississippi.	Stems.	1	.04		
<i>Balsamorhiza</i> sp.	do.	Texas.	Whole.	1	.23		
<i>Beaumontia grandiflora</i> (Roth) Wall.	Apocynaceae.	Utah.	Whole.	1	4.44		
<i>Barberris trifoliolata</i> Moric.	do.	Florida.	do.	1	1.25		
<i>Boerhaavia spicata</i> Choisy.	do.	Texas.	Leaves.	1	.09		
<i>Borreria frutescens</i> (L.) DC.	Nyctaginaceae.	do.	Roots.	1	0	.44	.22
<i>Brickellia dentata</i> (DC.) Sch. Bip.	Compositae.	Arizona.	Stems.	1	0		
<i>Bryophyllum crenatum</i> Baker.	do.	Texas.	Whole.	1	.37		
<i>Bumelia spinosa</i> A. DC.	Crassulaceae.	do.	do.	1	.09		
<i>texana</i> Buckl.	Sapotaceae.	South Carolina.	Leaves.	1	2.08		
<i>tenax</i> (L.) Willd.	do.	Texas.	Stems.	1	.04		
<i>Calotropis gigantea</i> (L.) R. Br.	do.	do.	Bark.	1	.75		
<i>procera</i> Ait.	do.	do.	Stems.	1	.10		
<i>Campanula persicifolia</i> L.	do.	Florida.	Bark.	1	.81		
sp.	do.	No record.	Leaves.	1	1.14		
<i>Calotropis gigantea</i> (L.) R. Br.	Asclepiadaceae.	do.	Leaves.	1	.99		
<i>procera</i> Ait.	do.	do.	Twigs.	1	.61		
<i>Campanula persicifolia</i> L.	Campanulaceae.	Oregon.	Leaves.	1	.78		
sp.	do.	do.	Stems.	1	.09		
<i>Canua</i> sp.	do.	Oregon.	Roots.	1	0		
<i>Canadus velutinus</i> Dougl.	Canaceae.	Maryland.	Leaves.	1	1.17		
	Sapindaceae.	Arkansas.	Stems.	1	.38		
		Oregon.	Leaves.	2	.08	.12	.10
			Stems.	2	.05	.10	.07

See footnote at end of table.

TABLE 1.—Rubber content of miscellaneous native and introduced plants collected in continental United States, analyzed 1920-45—Continued

Genus and species	Family	State where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Colastrus</i> sp.	Celastraceae	Maryland	{Leaves.....	1	Percent	Percent	Percent
			{Stems.....	1	.35		
<i>Conocaula americana</i> Nutt.	Compositae	Texas	{Roots.....	1	.34		
<i>Cornus</i> sp.	Cacaceae	California	{Whole.....	1	.03		
			{do.....	1	0		
<i>Clonitilla juavea</i> L.	Compositae	District of Columbia	{Stems.....	1	.40		
			{Roots.....	1	.51		
<i>Chondrophora virginica</i> (Nutt.) Greene	do.	Florida	{Whole.....	2	1.45	1.56	1.50
<i>Chrysopsis aspera</i> Shuttlw.	do.	South Carolina	{Roots.....	1	.31		
			{Whole.....	1	.30		
	do.	Maryland	{Leaves.....	1	2.84		
			{Stems.....	1	.34		
			{Whole.....	3	.39	3.14	2.21
			{Stems.....	3	.34	1.60	.86
			{Gum.....	2	22.43	29.45	25.96
			{Whole.....	4	42	1.44	90
			{Stems.....	5	.19	1.90	1.12
<i>Chrysoshamnus bigelovii</i> (Gray) Greene	do.	New Mexico	{Roots.....	1	1.08		
			{Whole.....	4	.20	1.26	.54
			{Stems.....	4	.86	.44	
			{Whole.....	2	.23	.48	.36
	do.	Wyoming	{Whole.....	1	.04		
			{Stems.....	1	1.71		
			{Branches.....	2	.16	.19	.18
			{Whole.....	2	.18	.36	.27
	do.	New Mexico	{Whole.....	0			
<i>Cichorium intybus</i> L.	do.	Maryland	{Whole.....	1			
<i>Cleome</i> sp.	Capparidaceae	No record	{do.....	1	.20		
<i>Commelina erecta</i> L.	Commelinaceae	Texas	{do.....	1	.21		
<i>Conyza collarii</i> Gray	Compositae	New Mexico	{do.....	1	.46		
<i>Croton monanthogynus</i> Michx.	Euphorbiaceae	Texas	{do.....	1	.03		
			{do.....	1	0		
			{do.....	1			
			{do.....	1	.04		
			{do.....	1	(¹)		
			{do.....	1	.11		
<i>Cytisus linifolius</i> Lam.	Leguminosae	California	{do.....	1	.12		
<i>Dioscorea batatas</i> Decne.	Dioscoriaceae	Georgia	{Tubers.....	1	.11		
			{do.....	1	.09		
			{do.....	2	0	.03	.02
			{do.....	1	0		
<i>Diospyros</i> sp.	Ebenaceae	Kentucky, Virginia	{Gum.....	1	.32		
			{Stems.....	1	.10		
			{Whole.....	1	.33		
<i>Dirca palustris</i> L.	Thymelaeaceae	Arkansas, Virginia	{Leaves.....	2	.12	.18	.15
<i>Echysanthera utilis</i> Hay. & Kaw.	Apocynaceae	Florida	{Whole.....	2			

TABLE 1.—Rubber content of miscellaneous native and introduced plants collected in continental United States, analyzed 1920-45—Continued

Genus and species	Family	State where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Euphorbia ammannioides</i> H. B. K.—Continued							
<i>heterophylla</i> L.	Euphorbiaceae	Arkansas, Florida, Mississippi	Leaves	1	Percent .42	Percent .42	Percent .26
<i>hypsipifolia</i> L.	do.	Texas	Stems	2	.11	.42	
<i>laeta</i> Engelm.	do.	do.	Roots	1	.07		
<i>lathyris</i> L.	do.	do.	Whole	1	.77		
<i>maculata</i> L.	do.	do.	do.	1	.21		
<i>marginata</i> Pursh.	do.	do.	Leaves	3	.09		
<i>mauretanica</i> L.	do.	California, Oregon, Pennsylvania, Washington, District of Columbia	Stems	3	.12	.30	.22
<i>misera</i> Benth.	do.	Oklahoma, Virginia	Roots	3	.23	.12	
<i>nutans</i> Lag.	do.	Nevada, Ohio, Texas	Whole	4	.01	.04	.02
<i>peplis</i> L.	do.	California	Whole	3	.08	.20	.12
<i>polychroma</i> Kern.	do.	do.	Whole	4	.09	.34	.21
<i>polyphylla</i> Engelm.	do.	Alabama	Leaves	2	.46	.54	.50
<i>prostrata</i> Alt.	do.	California	Stems	3	0	.18	.07
<i>pulcherrima</i> Willd.	do.	California, Florida, Texas	Roots	2	.02	.02	.02
<i>serpens</i> H.B.K.	do.	Texas	Gum	1	1.04		
<i>serrata</i> Engelm.	do.	do.	Whole	1	.35		
<i>stictospora</i> Engelm.	do.	do.	Stems	1	1.19		
<i>supina</i> Raf.	do.	do.	Roots	1	.14		
<i>titicaculi</i> L.	do.	do.	Whole	1	.27		
<i>virginiana</i> L.	do.	do.	Leaves	1	.29		
<i>virginiana</i> L.	do.	do.	Stems	1	.14		
<i>virginiana</i> L.	do.	do.	Whole	1	.21		
<i>virginiana</i> L.	do.	do.	Whole	1	0		
<i>virginiana</i> L.	do.	do.	Leaves	2	.18	.13	.12
<i>virginiana</i> L.	do.	do.	Stems	3	.09	.63	.43
<i>virginiana</i> L.	do.	do.	Lower	5	.02	.14	.08
<i>virginiana</i> L.	do.	do.	Middle	8	.02	.13	.07
<i>virginiana</i> L.	do.	do.	Upper	8	.06	.19	.11
<i>virginiana</i> L.	do.	do.	Twigs	8	.13	.39	.23
<i>virginiana</i> L.	do.	do.	Gum	1	.11		
<i>virginiana</i> L.	do.	do.	Whole	1	6.30		
<i>virginiana</i> L.	do.	do.	do.	2	.15	.22	.18
<i>virginiana</i> L.	do.	do.	do.	4	.23	.27	.25
<i>virginiana</i> L.	do.	do.	do.	2	.08	.10	.09
<i>virginiana</i> L.	do.	do.	do.	2	0		
<i>virginiana</i> L.	do.	do.	Stems	1	.39		
<i>virginiana</i> L.	do.	do.	Twigs	1	1.09		
<i>virginiana</i> L.	do.	do.	Branches	1	1.36		
<i>virginiana</i> L.	do.	do.	Gum	2	2.41	7.72	5.03
<i>virginiana</i> L.	do.	do.	Whole	1	.40		

<i>uriquhai</i> Torr. & Gray	do.	Texas	{Stems. {Roots.	1	0	17	1
<i>Ficus aurea</i> Nutt.	Moraceae	Florida	{Stems. {Leaves.	1	65	1	1
<i>bonghalensis</i> L.	do.	do.	Gum.	1	76.64	1	1
<i>capensis</i> Thunb.	do.	No record.	Whole.	1	18	1	1
<i>carica</i> L.	do.	Texas	do.	1	01	1	1
<i>heterophylla</i> L. f.	do.	California	Branch.	1	0	1	1
<i>hispida</i> L. f.	do.	No record.	{Leaves. {Fruit.	1	35 1.21	1	1
<i>rubiginosa</i> Desf.	do.	California	{Fruit. {Whole.	1	18 41	1	1
<i>Flavaria linearis</i> Lag.	Compositae	Florida	Bark.	1	13	1	1
<i>Fouquieria splendens</i> Engelm.	Fouquieriaceae	Arizona	Whole.	1	0	1	1
<i>Fucus</i> sp.	Fucaceae	Connecticut	{Stems. {Roots.	1	15	1	1
<i>Gilguthia</i> sp.	Leguminosae	California	do.	1	1	1	1
<i>Glottidium vesicarium</i> (Jacq.) Harper	Leguminosae	Mississippi	Stems.	1	1	1	1
<i>Guaphalium obtusifolium</i> L.	Compositae	South Carolina	Whole.	1	28	1	1
<i>harneyana</i> DC.	do.	do.	do.	2	30	1	1
<i>purpureum</i> L.	do.	Arkansas	do.	1	1.08	1	1
<i>Grindelia grandiflora</i> Hook.	do.	Texas	do.	1	40	1	1
<i>squarrosa</i> (Pursh) Dunal	do.	do.	{Stems. {Leaves.	1	3.57 57	1	1
sp.	do.	Montana	{Roots. {Flowers.	1	03 20	1	1
<i>Gutierrezia sarothrae</i> (Pursh) Britt. & Rusby	do.	California, Colorado	{Stems. {Leaves.	1	48 11	1	1
<i>Helentium</i> sp.	do.	Colorado	Whole.	1	14	1	1
<i>Heterodocha subaxillaris</i> (Lam.) Britt. & Rusby	do.	No record.	do.	1	0	1	1
<i>Hibiscus esculentus</i> L.	do.	South Carolina	do.	1	25	1	1
<i>Hosta ventricosa</i> (Salisb.) Stearn.	Malvaceae	Georgia	Pods.	1	0	1	1
<i>Hymenopappus scabosus</i> L'Herit.	Liliaceae	New York	Whole.	1	28	1	1
<i>Hymenoxys odorata</i> DC.	Compositae	Texas	do.	1	09	1	1
<i>richardsonii</i> (Hook.) Ckl.	do.	No record.	do.	1	16	1	1
<i>rudyi</i> (Gray) Ckl.	do.	New Mexico	{Stems. {Roots.	5	1.62 6.30	1.70 3.25	1
sp.	do.	Arizona	Whole.	1	28	1	1
<i>Ipomoea batatas</i> (L.) Lam.	do.	do.	do.	1	2	1	1
<i>Ira</i> sp.	Convolvulaceae	Florida	{Fruit. {Leaves.	1	05 57	1	1
<i>Jatropha berlandieri</i> Torr.	Anthusiaceae	No record.	Whole.	1	11	1	1
<i>multifida</i> L.	Euphorbiaceae	Texas	do.	1	52	1	1
<i>spathulata</i> (Orteg.) Muell. Arg.	do.	No record.	Leaves.	1	14	26	21
<i>texana</i> Muell. Arg.	do.	Texas	{Stems. {Roots.	3 2	08 11	10 20	1
<i>Jussiaea peruviana</i> L.	do.	Oklahoma, Texas	Whole.	2	13	27	1
sp.	Onagraceae	No record.	{Roots. {Whole.	2 1	05 10	.13 .09	1
<i>Jussiaea peruviana</i> L.	do.	do.	Leaves.	3	89	2.76	1
sp.	do.	Maryland	{Stems. {Roots.	1 1	09 09	1	1

TABLE 1.—*Rubber content of miscellaneous native and introduced plants collected in continental United States, analyzed 1920-45—Continued*

Genus and species	Family	State where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Kopsia arborea</i> Blume.	Apocynaceae.	No record.	Leaves.	10	Percent 2.96	Percent 6.81	Percent
<i>Lachnostoma arizonicum</i> Gray.	Asclepiadaceae.	Arizona.	Leaves.	1	2.96		
			Stems.	2	35	42	38
			Roots.	2	27	38	32
			Leaves.	3	11	68	43
<i>Lactuca canadensis</i> L.	Compositae.	Illinois, Maryland, Missouri, North Carolina, Oklahoma, Tennessee.	Stems.	3	12	41	25
			Whole.	4	4	33	19
			Gum.	1	4.36		
<i>floridana</i> (L.) Gaertn.	do.	Tennessee.	Leaves.	1	39		
			Stems.	1	06		
<i>indica</i> L.	do.	Maryland.	Leaves.	1	16		
<i>ludoviciana</i> (Nutt.) Ruddell.	do.	Texas.	Stems.	1	14		
<i>sativa</i> L.	do.	New Jersey, Utah.	Whole.	1	04		
			Gum.	1	26		
<i>serriola</i> L.	do.	Georgia, Nevada, New Mexico, Ohio, Oklahoma, Texas.	Whole.	1	24		
			Leaves.	4	34	59	42
			Stems.	4	01	15	08
			Roots.	3	01	05	04
			Whole.	2	22	28	25
			Gum.	1	4.86		
			Leaves.	1	24		
<i>spicata</i> (Lam.) Hitchc.	do.	New Jersey, Tennessee.	Stems.	1	02		
			Whole.	1	39		
			Leaves.	1	0		
sp.	do.	Connecticut, Pennsylvania, District of Columbia.	Gum.	1	0		
			Stems.	1	08		
			Whole.	1	20		
sp. (celtuce)	do.	North Carolina, Pennsylvania.	Whole.	3	13	28	18
<i>Larrea divaricata</i> Cav.	Zygophyllaceae.	Texas.	Whole.	1	20		
<i>Leucophyllum</i> Benth.	Scrophulariaceae.	Maryland.	do.	1	05		
<i>Liquidambar</i> (L.) Willd.	Compositae.	Virginia.	do.	1	17		
<i>Lobelia cardinalis</i> L.	Hamamelidaceae.	Maryland.	Wood.	1	0		
<i>elongata</i> Small.	Lobeliaceae.	District of Columbia.	Whole.	1	41		
	do.		do.	1	31		
<i>Lygodesmia grandiflora</i> Torr. & Gray	Compositae.	Colorado, New Mexico.	Roots.	1	5.47		
<i>juncosa</i> D. Don.	do.	South Dakota.	Whole.	1	64		
<i>texana</i> (Torr. & Gray) Greene.	do.	Texas.	Whole.	1	06		
sp.	do.	Montana.	Whole.	1	09		
			Whole.	1	19		

<i>Maclura pomifera</i> (Raf.) Schneid.....	Moraceae.....	Kansas, Louisiana, Virginia.....	Fruit.....	3	13	.30	.23
<i>Macrocarystis pyriferia</i> (L.) C. Ag.....	Lamiaceae.....	California.....	Whole.....	1	.09		
<i>Madia glomerata</i> Hook.....	Compositae.....	Utah.....	Cooked.....	1	.07		
<i>Martynia louisiana</i> Mill.....	Martyniaceae.....	Maryland.....	do.....	1	.17		
<i>Matelea laevis</i> (Zucc.) Woodson.....	Asclepiadaceae.....	do.....	{Leaves.....	1	0		
<i>Meibomia sepiarium</i> (Nutt.) Vail.....	do.....	Florida.....	Whole.....	1	.35		
<i>Morus</i> sp.....	Moraceae.....	do.....	do.....	1	.97		
<i>Nelumbo acerifolia</i> Gaertn.....	Nymphaeaceae.....	Louisiana.....	Gum.....	1	1.48		
<i>Nerium oleander</i> L.....	Apocynaceae.....	Mississippi.....	Whole.....	1	.03		
<i>Nicotiana trigonophylla</i> Dunal.....	Solanaceae.....	Texas.....	{Leaves.....	2	.21	.57	.39
		Arizona.....	Stems.....	2	(1)	.27	.14
			Whole.....	1	.21		
			Leaves.....	2	.17	.69	.43
			Stems.....	4	0	.26	.08
<i>Onoseris biennis</i> L.....	Onagraceae.....	Michigan, North Carolina, New Hampshire.....	Branches.....	1	.01		.01
			Roots.....	2	0	.02	
			Leaves.....	2	.31		1.22
<i>Onosmodium virginianum</i> (L.) A. DC.....	do.....	North Carolina.....	Whole.....	1	.60	2.14	
		South Carolina.....	Leaves.....	1	.60		
<i>Opuntia</i> sp.....	Cactaceae.....	Kansas, Tennessee.....	{Stems.....	1	.02		
<i>Parlinsonia aculeata</i> L.....	Leguminosae.....	Texas.....	Whole.....	1	0		
<i>Parthenium incanum</i> H. B. K.....	Compositae.....	do.....	Juice.....	1	0		
			Whole.....	1	0		
		do.....	do.....	1	0		
<i>lyraum</i> Gray.....	do.....	do.....	Whole.....	1	.28		
<i>Pedilanthus macrocarpus</i> Benth.....	Euphorbiaceae.....	California.....	Leaves.....	2	.57	.83	.70
<i>pedifolius</i> Poir.....	do.....	No record.....	Stems.....	2	.64	.94	.79
<i>tithymaloides</i> Poir.....	do.....	do.....	Roots.....	2	.77	.81	.79
		do.....	do.....	1	.25		
		do.....	do.....	1	.23		
		do.....	do.....	1	.27		
<i>Peltandra virginica</i> (L.) Schott & Endl.....	Araceae.....	Arizona, California, Florida.....	{Twigs.....	1	.54		
		Florida.....	Whole.....	3	.44	.68	.53
<i>Periploca graeca</i> L.....	Asclepiadaceae.....	Maryland.....	Fruit.....	1	.08		
<i>Phaseolus vulgaris</i> L.....	Leguminosae.....	California.....	Leaves.....	1	1.15		
<i>Phytolacca americana</i> L.....	Phytolaccaceae.....	Pennsylvania.....	Hulls.....	1	.13		
<i>Plantago major</i> L.....	Plantaginaceae.....	North Carolina.....	Leaves.....	1	.28		
		Pennsylvania.....	Whole.....	1	.28		
		do.....	do.....	1	3.36		
<i>Pluchea fortida</i> (L.) DC.....	Compositae.....	Florida.....	Leaves.....	1	.16		
			Stems.....	1	.16		
<i>Portulaca oleracea</i> L.....	Portulacaceae.....	New York.....	Whole.....	1	.82		
		Michigan, West Virginia, Wisconsin.....	Inner bark.....	1	0		
<i>Prunus</i> sp.....	Rosaceae.....	do.....	Gum.....	1	0		
			Whole.....	1	.02		

See footnote at end of table.

TABLE 1.—Rubber content of miscellaneous native and introduced plants collected in continental United States, analyzed 1920-45—Continued

Genus and species	Family	State where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Pteridium aquilinum</i> (L.) Kuhn.....	Polypodiaceae....	Arizona, Idaho, Maryland, Oregon, Washington.	{Roots.....	6	Percent	Percent	Percent
<i>Pterocaulon pycnostachyum</i> (Michx.) Ell.	Compositae....	South Carolina.....	{Whole.....	2	.02	.06	.04
<i>Pyrrhopappus carolinianus</i> (Walt.) DC.	do.....	do.....	{Leaves.....	1	.06	.09	.08
<i>multicaulis</i> DC.	do.....	Oklahoma.....	{Stems.....	1	1.28		
<i>Rhus glabra</i> L.	Anacardiaceae	Texas.....	{Whole.....	1	.28		
<i>typhina</i> L.	do.....	do.....	{do.....	1	.15		
<i>Sabal louisiana</i> (Darby) Bomhard	Arecaceae.....	Missouri, New Mexico, Tennessee.	{Roots.....	1	.18		
<i>Salpiglossis</i> sp.	Solanaceae.....	Tennessee.....	{Whole.....	2	.08	.26	.17
<i>Sambucus coerules</i> Raf.	Adoxaceae.....	Texas.....	{Whole.....	1	.04		
<i>racemosa</i> L.	do.....	California.....	{Juice.....	1	0		
<i>sp.</i>	do.....	do.....	{Flowers.....	1	.24		
<i>Sandwina chamaecyparissus</i> L.	do.....	New York.....	{Leaves.....	1	.09		
<i>Sarcobatus vermiculatus</i> (Hook.) Torr.	Compositae.....	do.....	{Stems.....	1	1.45		
<i>Scorzonera hispanica</i> L.	Chenopodiaceae.....	do.....	{Leaves.....	1	.06		
<i>Socum</i> sp.	Compositae.....	Connecticut, Pennsylvania.....	{Berries.....	1	.33		
<i>Sedum</i> sp.	do.....	Maryland, Pennsylvania.....	{Stems.....	1	.06		
<i>Sesuvio ambrosioides</i> Rydb.	do.....	Colorado.....	{Whole.....	2	(¹) 15	5.06	
<i>smallii</i> Britton.....	do.....	do.....	{Stems.....	2	.31		.23
<i>tridenticalatus</i> Rydb.	do.....	do.....	{Leaves.....	3	.07		.06
<i>Smilax laurifolia</i> L.	do.....	District of Columbia.....	{Roots.....	1	.15		
<i>roundifolia</i> L.	do.....	do.....	{Whole.....	1	.16		
<i>sp.</i>	do.....	do.....	{Leaves.....	1	.20		
<i>Solidago altissima</i> L.	do.....	do.....	{Stems.....	1	.06		
<i>angustifolia</i> Ell.	do.....	do.....	{Leaves.....	1	.04		
<i>tridenticalatus</i> Rydb.	do.....	do.....	{Roots.....	1	.51		
<i>Smilax laurifolia</i> L.	do.....	do.....	{Stems.....	1	.13		
<i>roundifolia</i> L.	do.....	do.....	{Whole.....	1	.22		
<i>sp.</i>	do.....	do.....	{Fruit.....	1	.07		
<i>Solidago altissima</i> L.	do.....	do.....	{Seed coating.....	4	.16	.64	.32
<i>angustifolia</i> Ell.	do.....	do.....	{Whole.....	2	.50	3.46	1.98
<i>tridenticalatus</i> Rydb.	do.....	do.....	{Fruit.....	2	.39		
<i>Smilax laurifolia</i> L.	do.....	do.....	{Leaves.....	1	(¹) 1.19		
<i>roundifolia</i> L.	do.....	do.....	{Stems.....	12	1.38	6.34	3.37
<i>sp.</i>	do.....	do.....	{Roots.....	12	.20	.03	.10
<i>Solidago altissima</i> L.	do.....	do.....	{Whole.....	14	.03	.09	.06
<i>angustifolia</i> Ell.	do.....	do.....	{Whole.....	1	.42	1.42	.62

<i>arguta</i> Ait.	do.	New Hampshire, Vermont, Virginia	Leaves.	1.02	1
			Stems.21	1
			Whole.08	5
<i>bicolor</i> L.	do.	Maryland, South Carolina, Virginia	Leaves.	1.60	61
			Stems.89	3
			Whole.	1.20	1.03
<i>cucuta</i> L.	do.	Maryland.	Leaves.25	12
			Stems.23	.87
<i>culicoides</i> Torr.	do.	New Hampshire.	Leaves.	2.46	1
<i>canadensis</i> L.	do.	Maryland, New Hampshire, New York, Ver- mont.	Stems.33	1
<i>chapmanii</i> Gray.	do.	South Carolina.	Whole.86	2
<i>elongata</i> Nutt.	do.	California.	Whole.	2.00	1.43
<i>erecta</i> Pursh.	do.	Virginia.	do.44	.57
			Leaves.37	
<i>fistulosa</i> Mill.	do.	Florida, Maryland, South Carolina.	Whole.	3.11	1
			Leaves.	1.02	1
<i>flexicaulis</i> L.	do.	Maryland.	Stems.80	1
			Whole.01	3
<i>graminifolia</i> (L.) Salisb.	do.	Maryland, New Hampshire, New York, South Carolina, District of Columbia.	Leaves.	2.08	4
			Stems.	4.48	4
<i>juncea</i> Ait.	do.	Maryland, New Hampshire, Virginia.	Whole.	1.65	.81
<i>macrophylla</i> Pursh.	do.	New Hampshire, Vermont.	Leaves.	4.48	2.99
			Stems.08	.05
			Whole.	1.39	.88
<i>microcephala</i> (Greene) Bush.	do.	Florida, South Carolina.	Leaves.	1.67	3.14
<i>monticola</i> Torr. & Gray.	do.	Virginia.	Stems.12	.21
<i>neglecta</i> Torr. & Gray.	do.	Maryland.	Whole.18	.79
			Leaves.74	.95
<i>memorialis</i> Ait.	do.	Maryland, New Hampshire.	Stems.	1.07	.14
			Whole.28	.14
<i>odora</i> Ait.	do.	Georgia, Maryland.	Leaves.	1.68	.77
<i>dioica</i> Nutt.	do.	New York.	Stems.34	1.16
<i>patriflorescens</i> (Michx.) Gray.	do.	Alabama, Florida.	Whole.93	.22
<i>patula</i> Ait.	do.	New York.	Stems.07	.57
			Whole.21	.57
			Leaves.16	.41
			Stems.32	
			Whole.	3.53	3.00
			Leaves.20	.10
			Stems.04	.08
			Whole.	1.01	.72
			Leaves.18	
			Stems.36	
			Whole.28	
			Leaves.98	1.42
			Stems.	2.80	2.54
			Whole.11	.18
			Leaves.50	.60
			Stems.	2.55	2.33
			Whole.08	.99
			Leaves.61	
			Stems.59	
			Whole.96	1.34
			Leaves.07	.04
			Stems.	1.67	
			Whole.87	

See footnote at end of table.

TABLE 1.—*Rubber content of miscellaneous native and introduced plants collected in continental United States, analyzed 1920-45—Continued*

Genus and species	Family	State where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Solidago altissima</i> L.—Continued							
<i>puberula</i> Nutt.....	Compositae.....	Maryland, Pennsylvania, Virginia..	Leaves.....	6	Percent .99	Percent 2.35	Percent 1.61
			Stems.....	5	.26	.30	.28
			Whole.....	5	.19	.37	.27
<i>racemosa</i> Greene.....	do.....	Virginia, South Carolina.....	Leaves.....	4	.75	1.72	1.26
<i>randii</i> (Porter) Britt.....	do.....	Vermont.....	Stems.....	1	.06
<i>rigida</i> L.....	do.....	District of Columbia..	Whole.....	1	.27
<i>rugosa</i> Mill.....	do.....	District of Columbia..	Leaves.....	1	3.00	3.10	3.05
			Stems.....	2	.25
			Whole.....	1	.99
<i>sempervirens</i> L.....	do.....	Maryland, New Hampshire, New York, Virginia, District of Columbia.	Leaves.....	9	2.63	3.94	3.14
			Stems.....	6	.01	.29	.19
			Whole.....	8	1.74	1.55	1.09
			Leaves.....	7	3.94	2.84
			Stems.....	3	.09	.37	.18
			Whole.....	4	.35	.91	.56
			Leaves.....	8	2.40	6.70	4.12
<i>gigantea</i> Ait.....	do.....	Maryland, New Hampshire, New York, Virginia.	Stems.....	6	.05	.27	.12
<i>sparsiflora</i> Gray.....	do.....	Arizona.....	Whole.....	1	.93	1.58	.89
<i>speciosa</i> Nutt.....	do.....	Maryland.....	Leaves.....	2	2.50	3.12	2.81
			Stems.....	2	.40	.34	.34
			Whole.....	2	.23	.76	.50
<i>squarrosa</i> Muhl.....	do.....	New Hampshire, Virginia.....	Leaves.....	4	.56	2.32	1.28
			Stems.....	1	.30
			Whole.....	1	.23
<i>stricta</i> Ait.....	do.....	Louisiana, South Carolina.	Leaves.....	3	1.59	2.87	2.64
			Stems.....	1	.10
			Whole.....	1	.12
<i>tenuifolia</i> Pursh.....	do.....	Maryland, South Carolina.	Leaves.....	1	.30	2.91	2.72
			Stems.....	2	2.50
			Whole.....	1	.09
			Stems.....	1	.02
			Whole.....	2	.58	.76	.67
<i>torifolia</i> Ell.....	do.....	Florida, South Carolina.	Leaves.....	1	2.95
<i>uliginosa</i> Nutt.....	do.....	Maine.....	Whole.....	2	1.26	1.75	1.50
<i>ulmifolia</i> Muhl.....	do.....	Maryland, Virginia.	Leaves.....	2	1.28	1.84	.86
			Stems.....	2	.04	.27	.16
<i>unilugata</i> (DC.) Porter.....	do.....	New York, Vermont	Whole.....	3	.88	.75	.82
			Leaves.....	6	.31	1.21	.59

<i>Sonchus oleraceus</i> L.	do.	Illinois, Minnesota.	Leaves.....	2	10	31	20
<i>asper</i> (L.) Hill.	do.	District of Columbia.	Stems.....	2	09	27	18
<i>oleraceus</i> L.	do.	Florida, New York, Texas.	Whole.....	2	38	1, 16	77
<i>Sophora secundiflora</i> Lag.	Leguminosae.	Texas.	Leaves.....	1	22		
<i>Stephanomeria virgata</i> Benth.	Compositae.	California.	Stems.....	1	50		
<i>Sillingtonia spatulata</i> (Muell. Arg.) Snull.	Euphorbiaceae.	No record.	Pods.....	3	07	42	29
<i>texasana</i> I. M. Johnston.	do.	Texas.	Leaves.....	1	23		
<i>trecitana</i> (Muell. Arg.) I. M. Johnston.	do.	do.	Stems.....	1	30		
<i>Symplocarpus foetidus</i> (L.) Nutt.	Araceae.	Pennsylvania.	Roots.....	1	10		
<i>paraxanthus</i> sp.	Compositae.	Maryland, New Jersey.	Juice.....	1	03		
<i>Thecetes grandiflora</i> .	Apocynaceae.	No record.	Whole.....	1	20		
<i>peruviana</i> (Pers.) Merr.	do.	Texas.	Whole.....	0	10		
<i>sp.</i>	do.	do.	do.	1	04		
<i>Tidestromia lanuginosa</i> (Nutt.) Standl.	Amaranthaceae.	do.	do.	1	09		
<i>Tradescantia canaliculata</i> Raf.	Commelinaceae.	do.	Roots.....	4	18	36	30
<i>foliosa</i> Small.	do.	do.	Whole.....	1	62		
<i>occidentalis</i> (Britton) Smyth.	do.	do.	Leaves.....	1	44		
<i>Tragopogon dubius</i> Scop.	Compositae.	Illinois, Indiana, Missouri, Ohio.	Stems.....	1	19		
<i>portifolius</i> L.	do.	Florida.	Whole.....	1	23		
<i>pratensis</i> L.	do.	Wisconsin.	Gum.....	1	57		
<i>sp.</i>	do.	Kansas, Montana.	Whole.....	0	30		
<i>Urtica esculenta</i> Benth.	Apocynaceae.	California, Nebraska, Nevada.	Stems.....	1	07		
<i>Varilla texana</i> Gray.	Compositae.	do.	Roots.....	3	26	59	46
<i>Verbesina occidentalis</i> (L.) Walt.	do.	do.	Whole.....	1	68		
<i>Vinca major</i> L.	Apocynaceae.	Michigan.	Leaves.....	1	14		
<i>Wyethia amplexicaulis</i> Nutt.	Compositae.	Oregon.	Stems.....	1	43		
<i>Zinnia</i> sp.	do.	do.	Roots.....	1	02		
		do.	Stems.....	1	18	23	20
		do.	Whole.....	2	32	45	38
		do.	Leaves.....	2	10	11	10
		do.	Roots.....	1	10		
		do.	Stems.....	1	35		
		do.	Whole.....	1	13		
		do.	Leaves.....	1	13		
		do.	Stems.....	1	4, 96		
		do.	Whole.....	1	13		
		do.	Leaves.....	1	96		
		do.	Stems.....	1	13		
		do.	Whole.....	1	09		
		do.	Leaves.....	1	1, 32		
		do.	Stems.....	2	62	1, 57	1, 10
		do.	Whole.....	2	17	22	20
		do.	Leaves.....	1	16		
		do.	Stems.....	1	04		
		do.	Whole.....	1	98		

1 Trace.

TABLE 2.—Total plants tested in table 1 and number containing rubber

Plants tested	Samples containing—			
	Rubber		No rubber	Total
	Above 0.10 percent	Below 0.10 percent		
Families.....	34	10	11	55
Genera.....	105	17	19	141
Species.....	240	34	29	303

TABLE 3.—List of plant families from table 1, showing number of species tested, number of samples analyzed, and number of samples containing rubber or no rubber

Family	Species	Tests	Samples containing—	
			Rubber ¹	No rubber ²
	Number	Number	Number	Number
Aceraceae.....	1	1	1	..
Adoxaceae.....	2	7	7	..
Aesculaceae.....	1	3	..	3
Amaranthaceae.....	1	1	..	1
Ambrosiaceae.....	1	1	1	..
Anacardiaceae.....	2	4	4	..
Apocynaceae.....	15	62	61	1
Araceae ³	2	2	2	..
Arecaceae ³	1	1	1	..
Aristolochiaceae.....	1	1	1	..
Asclepiadaceae.....	25	84	80	4
Berberidaceae.....	1	3	..	3
Boraginaceae.....	1	2	2	..
Cactaceae.....	2	3	..	3
Campanulaceae.....	2	6	4	2
Cannaceae.....	1	1	1	..
Capparidaceae.....	1	1	1	..
Celastraceae.....	4	8	8	..
Chenopodiaceae ³	1	3	3	..
Commelinaceae.....	4	8	8	..
Compositae.....	131	575	568	7
Convolvulaceae.....	1	1	1	..
Crassulaceae.....	2	2	1	1
Dioscoriaceae.....	3	4	3	1
Ebenaceae.....	2	2	..	1
Equisetaceae ³	1	1	1	..
Eucommiaceae.....	1	8	8	..
Euphorbiaceae.....	44	144	135	9
Fouquieriaceae.....	1	1	1	..
Fucaceae.....	2	2	..	2
Hamamelidaceae.....	1	1	..	1
Laminariaceae ³	1	1
Leguminosae.....	7	14	12	2
Liliaceae.....	2	2	2	..
Lobeliaceae.....	2	2	2	..
Malvaceae.....	1	1	..	1
Martyniaceae.....	1	2	2	..
Moraceae.....	9	15	14	1
Nyctaginaceae.....	1	1	1	..
Nymphaeaceae ³	1	1	1	..
Onagraceae.....	3	20	18	2
Phytolaccaceae.....	1	1	1	..
Plantaginaceae.....	1	2	2	..
Polypodiaceae ³	1	8	8	..
Portulacaceae.....	1	1	..	1
Rhodophyceae.....	1	1	..	1
Rosaceae ³	1	3	2	1
Rubiaceae.....	1	2	2	..
Sapindaceae.....	1	4	4	..
Sapotaceae.....	3	6	6	..
Scrophulariaceae ³	1	1	1	..
Simaroubaceae.....	1	2	2	..
Solanaceae.....	2	2	1	1
Thymelaeaceae.....	1	2	2	..
Zygophyllaceae.....	1	1	1	..

¹ A trace or more rubber found in at least 1 sample.² No rubber found in any sample.³ No analysis higher than 0.10 percent.

TABLE 4.—Rubber content of miscellaneous plants collected from outside continental United States

Genus and species	Family	Place where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Artocarpus</i> sp.	Moraceae	Haiti, P. R.	Latex.	2	Percent	Percent	Percent
<i>Bumelia lucidifera</i> HBK.	Sapotaceae	Sonora, Mexico	Fruit	1	3.54	3.83	3.68
<i>Casadipluvia cocaluco</i> Humb. & Bonpl.	Leguminosae	Sinaloa, Mexico	(Pod)	1	4.70
<i>Calatropis</i> sp.	Asclepiadaceae	Venezuela	(Stem)	1	15
<i>Camararia oblongifolia</i> Britton	Apocynaceae	Venezuela	Leaves	1	78
<i>Carissa grandiflora</i> A.DC.do.....	Puerto Rico	Stems	1	14
<i>Castilla elastica</i> Cav.	Moraceae	No record	Twigs	1	77
<i>Eclipta</i> sp.	Apocynaceae	Mexico	Stems	1	34
<i>Elaeophorbia drupifera</i> (Thonn.) Stapf.do.....	No record	Whole	2	2.08	2.28	2.18
<i>Euphorbia antisiphilitica</i> Zucc.do.....	Mexico	(Bark)	6	36	.85	.54
<i>callataoides</i> Benth.do.....	Puerto Rico	(Trunk)	2	.09	.16	.12
<i>filiza</i> Stapf.do.....	Cuba	Whole	1	1.01
<i>heptagona</i> L.do.....do.....	Leaves	1	26
<i>lactea</i> Haw.do.....do.....	(Gum)	1	3.30
<i>lactiflora</i> Phil.do.....	Sonora, Mexico	(Gum)	1	12.58
<i>lancifolia</i> Schlecht.do.....	Jalisco, Mexico	Twigs	1	1.27
<i>laro</i> Drake.do.....	No record	Whole	1	18.32
<i>latifolia</i> C. A. Mey.do.....	Dominican Republic	(Gum)	1	37
<i>nerifolia</i> L.do.....do.....	(Gum)	1	6.14
<i>portulacaoides</i> L.do.....	Chile	(Bark)	1	0
<i>schlechtendalii</i> Boiss.do.....	No record	(Whole)	1	11
<i>tirucalli</i> L.do.....	Cuba	(Gum)	2	8.34	9.00	8.67
sp.do.....	Saudi Arabia	Twigs	1	1.01
.....do.....do.....	(Gum)	2	4.90	5.62	5.26
.....do.....do.....	Whole	1	12
.....do.....do.....	(Gum)	2	2.05	2.27	2.16
.....do.....	Argentina	Twigs	1	22
.....do.....	Tamulipapas, Mexico	Roots	1	1.02
.....do.....	Cuba	Leaves	1	25
.....do.....do.....	Stems	1	1.04
.....do.....do.....	(Gum)	1	1.09
.....do.....do.....	Twigs	1	7.72
.....do.....do.....	(Bark)	1	0
.....do.....do.....	Twigs	1
.....do.....do.....	Outer	1

See footnote at end of table.

TABLE 4.—*Rubber content of miscellaneous plants collected from outside continental United States—Continued*

Genus and species	Family	Place where collected	Samples		Rubber content		
			Part	Number	Least	Most	Mean
<i>Ficus amara</i> Noronha.....	Moraceae.....	New Caledonia.....	Gum.....	1	Percent	Percent	Percent
<i>elastica</i> Roxb.....	do.....	{New Caledonia.....	do.....	1	7.12
<i>jimenezii</i> Standl.....	do.....	{Tamaulipas, Mexico.....	do.....	1	88.40
<i>petiolaris</i> H. B. K.....	do.....	{Guatemala.....	do.....	1	61.53	do.....
sp.....	do.....	{El Salvador.....	do.....	1	83.88	do.....
<i>Flourensia restiua</i> (Brandeg.) Blake.....	do.....	Sonora, Mexico.....	{Leaves.....	1	36
<i>Forsteronia portoricensis</i> Woodson.....	do.....	Chihuahua, Mexico.....	{Stems.....	1	44
<i>Hancornia speciosa</i> Gomez.....	Compositae.....	Ridalgo, Mexico.....	Gum.....	2	4.44	5.79	5.12
<i>Himatanthus articulatus</i> (Vahl) Woodson.....	Apocynaceae.....	Puerto Rico.....	Whole.....	1	56
<i>utropha albomaculata</i> Pax.....	do.....	Paraguay.....	do.....	1	24
sp. (chille).....	do.....	Paraguay.....	Gum.....	1	82.04
<i>Lucuma</i> sp.....	do.....	British Guiana.....	{Leaves.....	1	66
<i>Marsdenia edulis</i> S. Wats.....	do.....	Argentina.....	{Pods.....	1	70
<i>macrophylla</i> Fourn.....	Euphorbiaceae.....	Durango, Mexico.....	Whole.....	1	50
<i>verrucosa</i> Decne.....	do.....	Argentina.....	Gum.....	1	1.00
sp.....	do.....	{Jalisco, Mexico.....	do.....	2	14.58	30.95	22.76
<i>Mascarenhastia elastica</i> K. Schum.....	Sapotaceae.....	Sinaloa, Mexico.....	do.....	2	20.43	21.63	21.03
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Peru.....	do.....	9	19.74	33.74	26.83
<i>macrophylla</i> Fourn.....	do.....	Sonora, Mexico.....	{Leaves.....	1	92.26
<i>verrucosa</i> Decne.....	do.....	Tamaulipas, Mexico.....	{Stems.....	1	2.79
sp.....	do.....	Madagascar.....	{Leaves.....	1	54
<i>Mascarenhastia elastica</i> K. Schum.....	Apocynaceae.....	Sinaloa, Mexico.....	{Stems.....	1	3.39
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Cuba.....	Pod.....	1	80
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Nayarit, Mexico.....	Leaves.....	1	3.03
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Venezuela.....	Gum.....	1	3.40
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Dominican Republic.....	do.....	1	67.76
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Haiti.....	do.....	1	3.68
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Dominican Republic.....	{Leaves.....	1	37
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Sonora, Mexico.....	{Stems.....	1	1.04
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Venezuela.....	Fruit.....	1	85.74
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Dominican Republic.....	Gum.....	1	10.21
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Haiti.....	do.....	2	13.18	13.41	13.30
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Dominican Republic.....	do.....	2	13.58
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Sonora, Mexico.....	{Leaves.....	1	28
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Argentina.....	{Stems.....	1	60
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Nuevo Leon, Mexico.....	Whole.....	1	34
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Sinaloa, Mexico.....	Stems.....	1	40
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Tamaulipas, Mexico.....	{Leaves.....	1	73
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Mexico.....	{Stems.....	1	36
<i>Maidea lanata</i> (Zucc.) Woodson.....	do.....	Mexico.....	Gum.....	2	2.75	4.56	3.66

<i>mollis</i> H.B.K.....	do.....	Sonora, Mexico.....	(Fruit.....	1	36
		Mexico.....	Stems.....	1	61
<i>Sapium aucuparium</i> Jacq.....	Euphorbiaceae.....	Colombia.....	Gum.....	3	6.33	8.22
<i>biloculare</i> (S. Wats.) Pax.....	do.....	Argentina.....	Whole.....	1	17
		Sonora, Mexico.....	Stems.....	1	09
sp.....	do.....		Leaves.....	1	20
		Ecuador.....	Stems.....	3	07	.65
			Leaves.....	3	22	
<i>Scorzonera pulchra</i> Lomack.....	Compositae.....		Stems.....	2	21	.30
<i>Solidago chilensis</i> Meyen.....	do.....	Iraq.....	Bark.....	4	74	1.18
<i>Stemmadenia donnell-smithii</i> (Rose) Woodson.....	do.....	Argentina.....	(Gum.....	5	52.88	66.77
<i>palmeri</i> Rose & Standl.....	do.....	Honduras, Guatemala... ..	Roots.....	1	76	
		Sinaloa, Mexico.....	(Gum.....	1	11.57	
sp.....	do.....		Leaves.....	1	3.76	
		Guatemala.....	Stems.....	1	04	
<i>Thaetia ovata</i> A.DC.....	Apocynaceae.....		Gum.....	2	8.26	10.41
		Jalisco, Mexico... ..	Leaves.....	1	21	
			Twigs.....	1	3.21	
		Nayarit, Mexico.....	Whole.....	1	7.93	
			Fruit.....	1	86	
			Twigs.....	1	1	
			(Gum.....	2	13.58	16.51
<i>Vincetoxicum</i> sp.....	Asclepiadaceae.....	Guatemala.....	Leaves.....	1	10	
			Branches.....	1	54	
			Leaves.....	1	5.85	
			Branches.....	1	46	
			Fruit.....	1	34	
			(Fruit.....	1	50	
			Whole.....	1	27	

1 Trace.

TABLE 5.—*List of plant families from table 4, showing number of species tested, number of samples analyzed, and number of samples with and without rubber*

Family	Species	Tests	Samples containing—	
			Rubber	No rubber
	<i>Number</i>	<i>Number</i>	<i>Number</i>	<i>Number</i>
Apocynaceae.....	10	25	25
Asclepiadaceae.....	7	14	14
Compositae.....	7	17	17
Euphorbiaceae.....	19	61	59	2
Leguminosae.....	1	2	2
Moraceae.....	6	19	19
Sapotaceae.....	4	6	6

Conclusions

No new valuable rubber crop has been found. However, much information of scientific value has resulted. Through this extensive but uncoordinated survey more information is available about plants that accumulate rubber. Together with the surveys of Hall and Goodspeed (4), Hall and Long (5), Buehrer and Benson (1), Mitchell, Rice, and Roderick (8), Moxon and Whitehead (9), and the much more extensive but as yet unpublished work of Thomas G. Edison, this survey gives a comprehensive view of the plants in the United States that synthesize rubber.

Literature Cited

- (1) BEUHRER, T. F., and BENSON, L.
1945. RUBBER CONTENT OF NATIVE PLANTS OF THE SOUTHWESTERN DESERT. Ariz. Agr. Expt. Sta. Tech. Bul. 108, 33 pp.
- (2) COOK, O. F.
1903. THE CULTURE OF THE CENTRAL AMERICAN RUBBER TREE. U. S. Bur. Plant Indus. Bul. 49, 86 pp., illus.
- (3) FIGART, D. M.
1925. THE PLANTATION RUBBER INDUSTRY IN THE MIDDLE EAST. U. S. Bur. Foreign and Dom. Com. Trade Prom. Ser. 2, 317 pp., illus. [Crude Rubber Survey.]
- (4) HALL, H. M., and GOODSPEED, T. H.
1919. RUBBER PLANT SURVEY OF NORTHWESTERN NORTH AMERICA. Calif. Univ., Pubs., Bot. 7: 159-278, illus.
- (5) ——— and LONG, F. L.
1921. RUBBER CONTENT OF NORTH AMERICAN PLANTS. Carnegie Inst. Wash. Pub. 313, 65 pp., illus.
- (6) HOLT, E. G.
1927. MARKETING OF CRUDE RUBBER (WITH TRADE STATISTICS). U. S. Bur. Foreign and Dom. Com. Trade Prom. Ser. 55, 257 pp., illus. [Crude Rubber Survey.]
- (7) LA RUE, C. D.
1926. THE HEVEA RUBBER TREE IN THE AMAZON VALLEY. U. S. Dept. Agr. Dept. Bul. 1422, 69 pp., illus.
- (8) MITCHELL, J. H., RICE, M. A., and RODERICK, D. B.
1942. RUBBER ANALYSIS OF PLANTS IN SOUTH CAROLINA. Science 95: 624-625.

- (9) MOXON, A. L., and WHITEHEAD, E. I.
1943. THE RUBBER CONTENT OF SOME SOUTH DAKOTA PLANTS. S. Dak. Acad. Sci. Proc. 23: 53-55.
- (10) POLHAMUS, L. G.
1928. EXPERIMENTAL TAPPING OF HEVEA RUBBER TREES AT BAYEUX, HAITI, 1924-25. U. S. Dept. Agr. Tech. Bul. 65, 31 pp., illus.
- (11) ———
1933. RUBBER CONTENT OF VARIOUS SPECIES OF GOLDENROD. Jour. Agr. Res. 47: 149-152.
- (12) RANDS, R. D.
1924. SOUTH AMERICAN LEAF DISEASE OF PARA RUBBER. U. S. Dept. Agr. Dept. Bul. 1286, 18 pp., illus.
- (13) SCHURZ, W. L., HARGIS, O. D., MARBUT, C. F., and MANIFOLD, C. B.
1925. RUBBER PRODUCTION IN THE AMAZON VALLEY. U. S. Bur. Foreign and Dom. Com. Trade Prom. Ser. 23, 369 pp., illus. [Crude Rubber Survey.]
- (14) TREADWELL, J. C., HILL, C. R., and BENNETT, H. H.
1926. POSSIBILITIES FOR PARA RUBBER PRODUCTION IN NORTHERN TROPICAL AMERICA. U. S. Bur. Foreign and Dom. Com. Trade Prom. Ser. 40, 375 pp., illus. [Crude Rubber Survey.]
- (15) VANCE, C. F., MUZZALL, A. H., BUSHNELL, J. P., and BALDWIN, M.
1925. POSSIBILITIES FOR PARA RUBBER PRODUCTION IN THE PHILIPPINE ISLANDS. U. S. Bur. Foreign and Dom. Com. Trade Prom. Ser. 17, 101 pp., illus. [Crude Rubber Survey.]
- (16) VANDER LAAN, J. W.
1927. PRODUCTION OF GUTTA-PERCHA, BALATA, CHICLE AND ALLIED GUMS. U. S. Bur. Foreign and Dom. Com. Trade Prom. Ser. 41, 72 pp., illus. [Crude Rubber Survey.]
- (17) WEIR, J. R.
1926. A PATHOLOGICAL SURVEY OF THE PARA RUBBER TREE (HEVEA BRASILIENSIS) IN THE AMAZON VALLEY. U. S. Dept. Agr. Dept. Bul. 1380, 129 pp., illus.
- (18) WHITFORD, H. N., and ANTHONY, A.
1926. RUBBER PRODUCTION IN AFRICA. U. S. Bur. Foreign and Dom. Com. Trade Prom. Ser. 34, 136 pp., illus.

NATIONAL AGRICULTURAL LIBRARY



1022430350

NATIONAL AGRICULTURAL LIBRARY



1022430350